

Figure 1: PURRtentio a) DIY electrochemical biosensor b) Litter Box System and mobile application readouts. c) Components.

ABSTRACT

Traditional collection and analysis of feline urine samples for health monitoring are invasive, expensive, and infrequent. This paper introduces PURRtentio, a novel litter box system utilizing an electrochemical biosensor to monitor analytes in feline urine. The system comprises a DIY biosensor, potentiostat, microcontroller, distance sensor, and mobile application. Performance validation compared PURRtentio with an industry-grade potentiostat. PURRtentio presents an innovative and non-invasive approach for consistent monitoring of chemistry elements in feline urine, enabling early detection and management of cat's health conditions. This technology has the potential to revolutionize feline health monitoring, providing a solution for veterinarians and pet owners.

CCS CONCEPTS

• Human-centered computing \rightarrow Interactive systems and tools; Interaction devices; • Applied computing \rightarrow Life and medical sciences.

ACM ISBN 979-8-4007-0096-5/23/10.

KEYWORDS

biosensor, electrochemistry, potentiostat, urinalysis

ACM Reference Format:

Shuyi Sun, Gabriela Vega, Erkin Şeker, Krystle Reagan, and Katia Vega. 2023. PURRtentio: Implementing a Smart Litter Box for Feline Urinalysis with Electrochemical Biosensors. In The 36th Annual ACM Symposium on User Interface Software and Technology (UIST '23 Adjunct), October 29-November 1, 2023, San Francisco, CA, USA. ACM, New York, NY, USA, 3 pages. https:// //doi.org/10.1145/3586182.3615820

1 INTRODUCTION AND RELATED WORKS

Electrochemical biosensors are devices that detect and measure specific analytes, such as glucose, sodium, pH, and metal ions, by utilizing electrochemical reactions at electrodes. Typically, they consist of a working electrode, a counter electrode, and a reference electrode, with a potentiostat controlling the electrical potential and current between them. These biosensors have shown promise in health monitoring applications, from diabetes management to drug efficacy assessment [2, 24, 25].

Feline urine analysis is crucial for assessing the health status of cats, providing insights into hydration levels, kidney function, and metabolic disorders [3, 5, 18]. Sodium, potassium, glucose, and pH are common analytes in cat urine and serve as indicators of underlying health conditions [4, 10, 22, 26]. As a proof of concept, we focus on sodium levels due to they can a) vary significantly by dietary intake, making it easy to detect fluctuations even in

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).

UIST '23 Adjunct, October 29-November 1, 2023, San Francisco, CA, USA

^{© 2023} Copyright held by the owner/author(s).

https://doi.org/10.1145/3586182.3615820

healthy cats [4], b) be an indicative marker of feline health [14, 16],
and c) be more useful as a marker when assessed in combination
with other anayltes, which follows our project goal of multiple
analytes analysis in the future. Previous research explored colorimetric biosensors and visual chemical indicators in cat litter for
detection purposes [5, 10, 26]. However, colorimetric sensors have
limitations in lifespan and reversibility compared to electrochemical
biosensors, making them impractical for continuous use [1, 13, 15].

125 This project introduces PURRtentio, a litter box system that inte-126 grates electrochemical biosensors for unobtrusive and continuous monitoring of feline health. We developed a cost-effective solution 127 using DIY electrodes and a low-cost potentiostat to promote hard-128 129 ware and software replicability. Our focus is on detecting sodium levels in cat urine as an initial step towards comprehensive health 130 monitoring. By enabling continuous monitoring of feline's health, 131 PURRtentio facilitates early detection of health issues and person-132 alized care without costly and time-consuming lab tests [17, 18]. 133

2 IMPLEMENTATION

134

135

162

163

164

165

166

167

168

174

We modified a two-layer litter box to house the hardware. The 136 bottom layer includes the potentiostat system (Adafruit Feather 137 138 M4 Express board, Adafruit Bluefruit LE SPI Friend module, and 139 1000uA Rodeostat Featherwing V0.3 R1) and a pee pad for fluid absorption. The biosensor receives fluids through a sift surface 140 topped with non-absorbent cat litter. A second funnel layer ensures 141 all liquids are directed to an opening to the biosensor. The upper 142 layer incorporates VL53L0X Distance sensor (ToF) that triggers 143 the system upon detecting the cat's presence and departure. See 144 Figure 1b. The system operates with Open Circuit Potential (OCP). 145 OCP refers to the voltage present when a circuit's terminals are 146 disconnected. The form factor is designed to facilitate urine flow, 147 148 allowing sufficient biosensor stabilization time before testing.

The electrochemical biosensor is made using antioxidant-free, 149 oxidation-resistant copper tape with an insulating adhesive, en-150 151 suring resistance to oxidation and corrosion [7, 12]. This material choice provides a cost-effective alternative to traditional materials for biosensors [8, 9, 11, 20]. Fabrication involves cutting the copper 153 tape with a home-crafter cutting machine to create electrodes with 154 155 dimensions of 0.7 inches wide by 2 inches long (Fig. 1a), selected based on fluid flow areas, flexibility, and testing requirements. The 156 working electrode undergoes treatment with a sodium activation so-157 lution, making it sodium ion-selective and preventing interference 158 159 from other elements [19]. An insulating barrier solution is applied to the exposed areas of the electrodes, not on the biosensor section, 160 161 ensuring precise measurements and prevent contamination.

The system operates asynchronously with optimized sampling rate to accommodate biosensor behavior and BLE communication with the application. Utilizing the p5ble.js library, the application establishes a connection with the device, enabling real-time data visualization through live graphs and concentration level gauges.

3 TECHNICAL EVALUATION

In choosing biosensor material, we compared our potentiostat in
PURRtentio with the industry-grade potentiostat EmStat. Sodium
test solutions were prepared using preferred electrochemical techniques [6, 21] at concentrations based on feline urine levels [4, 14,
23]: 43.48, 97.83, 156.52, 195.65, 234.8, and 391.3 mM. We prepared



Figure 2: Comparision of EmStat and RodeoStat

them through dilution of a highly concentrated of sodium solution with known tested value in deionized water. Solutions were mixed by machine controlling speed, time, temperature, and amount [21].

We conducted a technical evaluation to validate the performance of the DIY biosensor and the PURRtentio system by comparing them against EmStat using OCP with prepared test solutions. Our system show similar trends for either system used. This is evident from the trendline analyses and the r-squared values (shown in Figure 2), with slope values similar to the ten thousandth place and r-squared values <0.1 apart, which indicate the strength of the correlation between the measured values and the expected values.

4 CONCLUSION AND FUTURE WORKS

Our study presents PURRtentio, an innovative method for continuous and non-invasive monitoring of feline health using electrochemical biosensors. By developing a DIY biosensor and low-cost potentiostat, we accurately measure sodium levels in feline urine. Our evaluation shows that PURRtentio performs comparably to an industry-grade potentiostat, providing reliable measurements for different sodium concentrations.

Future work includes developing a rinsing system to prolong biosensor lifespan, conducting tests with various analytes and feline urine, interviewing veterinarians, and performing user studies in real-world settings. Integrating PURRtentio with IoT devices like smart feeders and activity trackers enhances functionality for comprehensive feline health monitoring. Exploring other analytes such as glucose for feline diabetes and simultaneous detection of multiple analytes holds great potential.

PURRtentio revolutionizes feline health monitoring, improving veterinary care and cat well-being. With further advancements and IoT integration, this technology has broader applications in animal and human health monitoring and beyond.

ACKNOWLEDGMENTS

This material is based upon work partially supported by the National Science Foundation under Grant No 2146461.

175

176

PURRtentio: Implementing a Smart Litter Box for Feline Urinalysis with Electrochemical Biosensors UIST '23 Adjunct, October 29-November 1, 2023, San Francisco, CA, USA

233 **REFERENCES**

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

273

274

275

276

277

278

279

280

281

282

283

284

285

290

- [1] Wilson A. Ameku, Josué M. Gonçalves, Vanessa N. Ataide, Mauro S. Ferreira Santos, Ivano G. R. Gutz, Koiti Araki, and Thiago R. L. C. Paixão. 2021. Combined Colorimetric and Electrochemical Measurement Paper-Based Device for Chemometric Proof-of-Concept Analysis of Cocaine Samples. ACS Omega 6, 1 (12 Jan 2021), 594–605. https://doi.org/10.1021/acsomega.0c05077
- [2] Ananta Narayanan Balaji, Chen Yuan, Bo Wang, Li-Shiuan Peh, and Huilin Shao. 2019. pH Watch-Leveraging pulse oximeters in existing wearables for reusable, real-time monitoring of pH in sweat. In Proceedings of the 17th Annual International Conference on Mobile Systems, Applications, and Services. 262–274.
- [3] Catherine Bovens. 2011. Feline Lower Urinary Tract Disease A diagnostic approach. Feline Update (2011).
- [4] Y. H. Cottam, P. Caley, S. Wamberg, and W. H. Hendriks. 2002. Feline Reference Values for Urine Composition. The Journal of Nutrition 132, 6 (06 2002), 1754S-1756S. https://doi.org/10.1093/jn/132.6.1754S arXiv:https://academic.oup.com/jn/article-pdf/132/6/1754S/30067020/1754s.pdf
- [5] Lanny U. Franklin and Mary E. Sachs. 1993. PH-INDICATING MATERAL AND CAT LTTER CONTAINING SAME.
- [6] Guinevere Giffin. 2022. The role of concentration in electrolyte solutions for non-aqueous lithium-based batteries. *Nature Communications* 13 (09 2022). https://doi.org/10.1038/s41467-022-32794-z
- [7] Alireza Khoshroo, Komail Sadrjavadi, Mojtaba Taran, and Ali Fattahi. 2020. Electrochemical system designed on a copper tape platform as a nonenzymatic glucose sensor. Sensors and Actuators B: Chemical 325 (08 2020), 128778. https: //doi.org/10.1016/j.snb.2020.128778
- [8] P. Khullar, J. Badilla, and R. Kelly. 2015. The Use of a Sintered Ag/AgCl Electrode as Both Reference and Counter Electrode for Electrochemical Measurements in Thin Film Electrolytes. ECS Electrochemistry Letters 4 (01 2015), C31–C33. https://doi.org/10.1149/2.0051510eel
- [9] Tae Yong Kim, Sung A Hong, and Sung Yang. 2015. A Solid-State Thin-Film Ag/AgCl Reference Electrode Coated with Graphene Oxide and Its Use in a pH Sensor. Sensors 15, 3 (2015), 6469–6482. https://doi.org/10.3390/s150306469
- [10] Charles R. Kuhns. 1992. ANIMAL LITTER WITH CHEMICALLY BOUND CHEM-ICAL INDICATORS.
- [11] Min Suk Lee, Akshay Paul, Yuchen Xu, W. David Hairston, and Gert Cauwenberghs. 2022. Characterization of Ag/AgCl Dry Electrodes for Wearable Electrophysiological Sensing. Frontiers in Electronics 2 (2022). https://doi.org/10.3389/ felec.2021.700363
- [12] Emil R. Mamleyev, Peter G. Weidler, Alexei Nefedov, Dorothée Vinga Szabó, Monsur Islam, Dario Mager, and Jan G. Korvink. 2021. Nano- and Microstructured Copper/Copper Oxide Composites on Laser-Induced Carbon for Enzyme-Free Glucose Sensors. ACS Applied Nano Materials 4, 12 (2021), 13747–13760. https: //doi.org/10.1021/acsanm.1c03149 arXiv:https://doi.org/10.1021/acsanm.1c03149
- [13] Keiichiro MIE, Akiyoshi HAYASHI, Hidetaka Nishida, Mari OKAMOTO, Kazuo YASUDA, Mio NAKATA, Kazuyuki FUKATSU, Norie MATSUNAMI, Shogo YA-MASHITA, Fumihito OHASHI, and Hideo AKIYOSHI. 2019. Evaluation of the accuracy of urine analyzers in dogs and cats. *Journal of Veterinary Medical Science* 81 (10 2019). https://doi.org/10.1292/jvms.18-0468
- [14] P. Nguyen, B. Reynolds, J. Zentek, N. Paßlack, and V. Leray. 2017. Sodium in feline nutrition. *Journal of Animal Physiology and Animal Nutrition* 101, 3 (2017), 403–420. https://doi.org/10.1111/jpn.12548 arXiv:https://onlinelibrary.wiley.com/doi/pdf/10.1111/jpn.12548
- [15] J. Pérez-Accino, L. Bernabe, E. Manzanilla, and J. Puig. 2020. The utility of combined urine dipstick analysis and specific gravity measurement to determine feline proteinuria. *Journal of Small Animal Practice* 61 (07 2020). https://doi.org/ 10.1111/jsap.13184
- [16] Yann Queau, Esther Bijsmans, Alexandre Feugier, and Vincent Biourge. 2020. Increasing dietary sodium chloride promotes urine dilution and decreases struvite and calcium oxalate relative supersaturation in healthy dogs and cats. *Journal of Animal Physiology and Animal Nutrition* 104 (03 2020). https://doi.org/10.1111/ jpn.13329
- [17] Rose E. Raskin, Kelly A. Murray, and Julie K. Levy. 2002. Comparison of Home Monitoring Methods for Feline Urine pH Measurement. Veterinary Clinical Pathology 31, 2 (2002), 51–55. https://doi.org/10.1111/j.1939-165X. 2002.tb00279.x arXiv:https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1939-165X.2002.tb00279.x
- [18] George Reppas and Susan F Foster. 2016. Practical urinalysis in the cat: 2: Urine microscopic examination 'tips and traps'. *Journal of Feline Medicine* and Surgery 18, 5 (2016), 373–385. https://doi.org/10.1177/1098612X16643249 arXiv:https://doi.org/10.1177/1098612X16643249 PMID: 27143040.
- [19] Blaine F. Severin and Thomas D. Hayes. 2021. Effect of electrode rinse solutions on the electrodialysis of concentrated salts. *Separation and Purification Technology* 274 (2021), 119048. https://doi.org/10.1016/j.seppur.2021.119048
- [26] [27] KK Mohammed Shafeeq, Shiny Nair, G Uma, and T Mukundan. 2019. Fabrication of Ag/AgCl electrode for detection of electric field in marine environment. In *IOP Conference Series: Materials Science and Engineering*, Vol. 561. IOP Publishing, 012054.

- [21] Nikola Slaninova, Klara Fiedorova, Ali Selamat, Karolina Danisova, Jan Kubicek, Ewaryst Tkacz, and Martin Augustynek. 2020. Analysis and Testing of a Suitable Compatible Electrode's Material for Continuous Measurement of Glucose Concentration. Sensors 20, 13 (2020). https://doi.org/10.3390/s20133666
- [22] Ralph J. Steckel and Larry J. Murphy. 2008. Animal litter having the property of detecting diabetes in felines.
- [23] Hui Xu, Dorothy P.L. Laflamme, and Grace L. Long. 2009. Effects of dietary sodium chloride on health parameters in mature cats. *Journal of Feline Medicine* and Surgery 11, 6 (2009), 435–441. https://doi.org/10.1016/j.jfms.2008.10.001 arXiv:https://doi.org/10.1016/j.jfms.2008.10.001 PMID: 19073369.
- [24] Yiran Yang and Wei Gao. 2019. Wearable and flexible electronics for continuous molecular monitoring. *Chemical Society Reviews* 48, 6 (2019), 1465–1491.
- [25] Meredith Young-Ng, Grace Chen, Danielle Kiyama, Anna-Sofia Giannicola, Erkin Şeker, and Katia Vega. 2023. Sweatcessory: a Wearable Necklace for Sensing Biological Data in Sweat. In Adjunct Proceedings of the 2022 ACM International Joint Conference on Pervasive and Ubiquitous Computing and the 2022 ACM International Symposium on Wearable Computers (UbiComp/ISWC '22 Adjunct). Association for Computing Machinery, New York, NY, USA, 141–143. https: //doi.org/10.1145/3544793.3560342
- [26] . 2019. Cat urine pH indicating material, pH indicating cat litter and preparation method thereof.

Received 19 July 2023

330

331

332

333

334

335

336

337

338

339

341

342

343

344

345

346

347

348

291

292

293

294